



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A61K 38/00, 49/00, C07K 5/00, 7/00, 17/00</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 95/22341</b> <b>(43) International Publication Date:</b> 24 August 1995 (24.08.95)
<b>(21) International Application Number:</b> PCT/US95/02131 <b>(22) International Filing Date:</b> 21 February 1995 (21.02.95) <b>(30) Priority Data:</b> 94200409.4 18 February 1994 (18.02.94) EP <b>(34) Countries for which the regional or international application was filed:</b> GB et al. <b>(71) Applicant (for all designated States except US):</b> MALLINCK-RODT MEDICAL, INC. [US/US]; 675 McDonnell Boulevard, P.O. Box 5840, St. Louis, MO 63134 (US). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> MERTENS, John [BE/BE]; Laarbeeklaan 103, B-1090 Brussels (BE). TOURWÉ, Dirk [BE/BE]; Pleinlaan 2, B-1050 Brussels (BE). CEUSTERS, Marc [BE/BE]; Pleinlaan 2, B-1050 Brussels (BE). <b>(74) Agents:</b> McBRIDE, Thomas, P. et al.; Mallinckrodt Medical, Inc., 675 McDonnell Boulevard, P.O. Box 5840, St. Louis, MO 63134 (US).		<b>(81) Designated States:</b> CA, HU, JP, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title: LABELLED PEPTIDE COMPOUNDS</b>		
<b>(57) Abstract</b>		
<p>The invention relates to a labelled peptide compound, wherein the peptide has a selective neurotensin receptor affinity and is represented by the general formula (I): <math>R_1 - (^1\text{Pro})_n - ^2\text{Xaa} - ^3\text{Xbb} - ^4\text{Xcc} - ^5\text{Xdd} - ^6\text{Xee} - ^7\text{Leu} - \text{OH}</math>, wherein <math>R_1</math> is a <math>(\text{C}_1\text{-C}_3)</math>alkanoyl group, an arylcarbonyl group, an aryl-<math>(\text{C}_1\text{-C}_3)</math>alkanoyl group, or a chelating group attached by an amide bond or through a spacing group to the peptide molecule; Xaa and Xbb are each individually Arg or Lys; Xcc is an unsubstituted or substituted cyclic amino acid, preferably selected from Pro and Hyp; Xdd is Tyr, Trp or Phe; Xee is Leu, Ile or t.-butylalanine; and <math>n</math> is 0 or 1; and wherein said peptide is labelled with (a) a metal isotope, or (b) with a radioactive halogen isotope; on the understanding that: (i) if the label is a metal isotope, <math>R_1</math> represents a chelating group for chelating said isotope; and (ii) if the label is a radioactive halogen isotope, said halogen isotope is attached to <math>^4\text{Tyr}</math> in the 2-position of the phenyl ring, to <math>^4\text{Trp}</math>, or to the aryl group of substituent <math>R_1</math>. The invention further relates to a pharmaceutical composition comprising said labelled peptide compound, to the use of said composition for diagnosis and therapy, and to a kit for preparing a radiopharmaceutical composition.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

Labelled peptide compounds

The present invention relates to labelled peptide compounds, to a method of preparing these compounds, to a pharmaceutical composition comprising these compounds, to the use of this composition for diagnosis and therapy, and to a kit for preparing a radiopharmaceutical composition.

Neurotensin is a thirteen amino acid peptide, in 1973 isolated from bovine hypothalamus. It has the following structure:

pGlu-Leu-Tyr-Glu-Asn-Lys-Pro-Arg-Arg-Pro-Tyr-Ile-Leu-OH

High concentrations of neurotensin are found in discrete regions of the mammalian central nervous system. In addition, neurotensin interacts with specific receptors in the periphery. During the last decade, neurotensin receptors were found in several tumour cells, like small cell lung carcinoma, human colon carcinoma and human meningiomas.

Two radioiodinated neurotensin derivatives are mentioned in the literature. Because of the existence of two tyrosine residues in the neurotensin sequence, iodination of neurotensin yields a complex mixture of radioiodinated peptides, that possess very different biological properties and, moreover, are difficult to purify. Therefore, Mazella et al. (J. Biol. Chem. 1983, 258, 3476-3481) synthesized a neurotensin analogue in which the tyrosine-11 is replaced by a tryptophan residue: [Trp<sup>11</sup>]neurotensin. Iodination (with 125-iodide under electrophilic conditions on the aromatic ring of tyrosine-3) results in a mono-iodo derivative showing a  $K_d$  of 0.1 nM for binding on rat brain synaptic membranes. Preliminary experiments from the same group of researchers (Sadoul et al.; Biochem. Biophys. Res. Commun. 1984, 120, 812-819) indicated that neurotensin receptors in human brain showed a low affinity for monoiodo-[Trp<sup>11</sup>]-neurotensin, making this compound unsuitable for binding experiments. The same group of researchers succeeded later on in preparing a mono-iodo derivative of neurotensin itself; only tyrosine-3 was iodinated with 125-iodide in this method. This radioiodinated analogue has a  $K_d$  of 0.2 nM for binding on rat brain

synaptic membranes and a  $K_d$  of 0.26 nM for binding to human brain neurotensin receptors.


The labelled natural neurotensin as well as the labelled tryptophan-11 neurotensin analogue, however, suffers from an enzymatic breakdown due to cleavage of peptide bonds, resulting in an in vivo half-life of only a few minutes.

Structure-activity studies (Granier et al.; Eur. J. Biochem. 1982, 124, 117-125) revealed that the right-hand part of the neurotensin molecule fulfills the structural requirements for mimicking the entire sequence, provided its  $\alpha$ -amino end group is protected by acetylation. The binding affinities of this analogue are comparable with those of neurotensin in two binding assays, viz. the binding assay on rat brain synaptic membranes and that on HT 29 cells.

This analogue contains one remaining tyrosine residue which can be readily radioiodinated using electrophilic substitution of the hydrogen in the ortho position of the phenolic group. Structure-activity studies revealed, however, that iodination in the 3-position of said tyrosine residue of this analogue results in a remarkable loss of receptor affinity, viz. with a factor of 20 (Mazella et al.; see above).

It is the objective of the present invention to provide a labelled peptide compound which has a selective affinity to neurotensin receptors, comparable with that of neurotensin itself, and which has a sufficient enzymatic resistance to allow its use in diagnosis and therapy.

This objective can be achieved by a labelled peptide compound, wherein the peptide has a selective neurotensin receptor affinity and is represented by the general formula



wherein:

$R_1$  is a  $(C_1-C_3)$ alkanoyl group, an arylcarbonyl group, an aryl- $(C_1-C_3)$ alkanoyl group, or a chelating group attached by an amide bond or through a spacing group to the peptide molecule;

Xaa and Xbb are each individually Arg or Lys;

Xcc is an unsubstituted or substituted cyclic amino acid, preferably selected from Pro and Hyp;

Xdd is Tyr, Trp or Phe;

Xee is Leu, Ile or t.-butylalanine; and

n is 0 or 1;

and wherein said peptide is labelled with (a) a metal isotope selected from the group consisting of <sup>99m</sup>Tc, <sup>203</sup>Pb, <sup>67</sup>Ga, <sup>68</sup>Ga, <sup>72</sup>As, <sup>111</sup>In, <sup>113m</sup>In, <sup>97</sup>Ru, <sup>62</sup>Cu, <sup>64</sup>Cu, <sup>52</sup>Fe, <sup>52m</sup>Mn, <sup>51</sup>Cr, <sup>186</sup>Re, <sup>188</sup>Re, <sup>77</sup>As, <sup>90</sup>Y, <sup>67</sup>Cu, <sup>169</sup>Er, <sup>121</sup>Sn, <sup>127</sup>Te, <sup>142</sup>Pr, <sup>143</sup>Pr, <sup>198</sup>Au, <sup>199</sup>Au, <sup>161</sup>Tb, <sup>109</sup>Pd, <sup>165</sup>Dy, <sup>149</sup>Pm, <sup>151</sup>Pm, <sup>153</sup>Sm, <sup>157</sup>Gd, <sup>159</sup>Gd, <sup>166</sup>Ho, <sup>172</sup>Tm, <sup>169</sup>Yb, <sup>175</sup>Yb, <sup>177</sup>Lu, <sup>105</sup>Rh and <sup>111</sup>Ag, or

(b) with a radioactive halogen isotope;

on the understanding that:

(i) if the label is a metal isotope, R<sub>1</sub> represents a chelating group for chelating said isotope; and

(ii) if the label is a radioactive halogen isotope, said halogen isotope is attached to 'Tyr in the 2-position of the phenyl ring, to 'Trp, or to the aryl group of substituent R<sub>1</sub>.


Suitable examples of aryl groups in R<sub>1</sub> are: phenyl, halo-substituted phenyl or indolyl; preferably phenyl, 4-fluorophenyl, 2- or 4-bromophenyl, 2-iodophenyl, 4-fluoro-2-bromophenyl and 4-fluoro-2-iodophenyl. Suitable examples of radioactive halogen isotopes are: <sup>123</sup>I, <sup>124</sup>I, <sup>125</sup>I, <sup>131</sup>I, <sup>75</sup>Br, <sup>76</sup>Br, <sup>77</sup>Br and <sup>82</sup>Br.

In the above labelled peptide compounds one or more of the amino acids may have the D-configuration instead of the normal L-configuration.

The labelled peptide compounds of the invention may also comprise so-called pseudo peptide bonds, viz. -CH<sub>2</sub>-NH- bonds, in addition to the natural amide bonds, viz. -CO-NH- bonds. Such modifications of the amino acids naturally occurring in peptides are within the scope of the present invention.

Peptide compounds which, according to the invention, are labelled with a metal isotope as indicated above, are provided, directly or through a spacing group, with a chelating group, said chelating group being attached by an amide bond to an amino group of said peptide compound.


Said chelating group is preferably derived from ethylene diamine tetra-acetic acid (EDTA), diethylene triamine penta-acetic acid (DTPA), cyclohexyl 1,2-diamine tetra-acetic acid (CDTA), ethyleneglycol-0,0'-bis(2-aminoethyl)-N,N,N',N'-tetra-acetic acid (EGTA), N,N-bis(hydroxybenzyl)-ethylenediamine-N,N'-diacetic acid (HBED), triethylene tetramine hexa-acetic acid (TTHA), 1,4,7,10-tetraazacyclododecane-N,N',N'',N'''-tetra-acetic acid (DOTA), hydroxyethylidiamine triacetic acid (HEDTA), 1,4,8,11-tetra-azacyclotetradecane-N,N',N'',N'''-tetra-acetic acid (TETA), substituted DTPA, substituted EDTA, or from a compound of the general formula



(III)

wherein R is a branched or non-branched, optionally substituted hydrocarbonyl radical, which may be interrupted by one or more heteroatoms selected from N, O and S and/or by one or more NH groups, and Y is a group which is capable of reacting with an amino group of the peptide and which is preferably selected from the group consisting of carbonyl, carbimidoyl, N-(C<sub>1</sub>-C<sub>6</sub>)alkylcarbimidoyl, N-hydroxycarbimidoyl and N-(C<sub>1</sub>-C<sub>6</sub>)alkoxycarbimidoyl.

Examples of suitable chelators of the general formula III are unsubstituted or substituted 2-iminothiolanes and 2-iminothiacyclohexanes, in particular 2-imino-4-mercaptomethylthiolane. Said optionally present spacing group has preferably the general formula



(IV)

(V)

wherein R<sub>3</sub> is a C<sub>1</sub>-C<sub>10</sub> alkylene group, a C<sub>1</sub>-C<sub>10</sub> alkylidene group or a C<sub>2</sub>-C<sub>10</sub> alkenylene group, and X is a thiocarbonyl group or a methylcarbonyl group.

In connection with their biological properties, in particular their binding affinity to neurotensin receptors and their in vivo stability, and with their synthetic accessibility, the following labelled peptide

compounds according to the invention are preferred:

- (1) (2-"I-phenyl)acetyl - <sup>1</sup>Arg - <sup>2</sup>Arg - <sup>3</sup>Pro - <sup>4</sup>Tyr - <sup>5</sup>Ile - <sup>6</sup>Leu - OH
- (2) (4-"I-phenyl)acetyl - <sup>1</sup>Arg - <sup>2</sup>Arg - <sup>3</sup>Pro - <sup>4</sup>Tyr - <sup>5</sup>Ile - <sup>6</sup>Leu - OH
- (3) 2-"I-benzoyl - <sup>1</sup>Arg - <sup>2</sup>Arg - <sup>3</sup>Pro - <sup>4</sup>Tyr - <sup>5</sup>Ile - <sup>6</sup>Leu - OH
- (4) 4-"I-benzoyl - <sup>1</sup>Arg - <sup>2</sup>Arg - <sup>3</sup>Pro - <sup>4</sup>Tyr - <sup>5</sup>Ile - <sup>6</sup>Leu - OH
- (5) (4-F-2-"I-phenyl)acetyl - <sup>1</sup>Arg - <sup>2</sup>Arg - <sup>3</sup>Pro - <sup>4</sup>Tyr - <sup>5</sup>Ile - <sup>6</sup>Leu - OH
- (6) (4-F-2-"I-benzoyl) - <sup>1</sup>Arg - <sup>2</sup>Arg - <sup>3</sup>Pro - <sup>4</sup>Tyr - <sup>5</sup>Ile - <sup>6</sup>Leu - OH

as well as variants of compounds (1) to (6), wherein <sup>1</sup>Arg is attached to <sup>2</sup>Arg by pseudo peptide bonds,

- (7) (2-"I-phenyl)acetyl - <sup>1</sup>Lys - <sup>2</sup>Arg - <sup>3</sup>Pro - <sup>4</sup>Tyr - <sup>5</sup>Ile - <sup>6</sup>Leu - OH
- (8) (2-"I-phenyl)acetyl - <sup>1</sup>Arg - <sup>2</sup>Lys - <sup>3</sup>Pro - <sup>4</sup>Tyr - <sup>5</sup>Ile - <sup>6</sup>Leu - OH
- (9) Ac - <sup>1</sup>Arg - <sup>2</sup>Arg - <sup>3</sup>Pro - <sup>4</sup>[2-"I-Tyr] - <sup>5</sup>Ile - <sup>6</sup>Leu - OH
- (10) Ac - <sup>1</sup>Arg - <sup>2</sup>Arg - <sup>3</sup>Pro - <sup>4</sup>[2-"I-Trp] - <sup>5</sup>Ile - <sup>6</sup>Leu - OH
- (11) Ac - <sup>1</sup>Arg - <sup>2</sup>Arg - <sup>3</sup>Pro - <sup>4</sup>[5-"I-Trp] - <sup>5</sup>Ile - <sup>6</sup>Leu - OH
- (12) Ac - <sup>1</sup>Arg - <sup>2</sup>Arg - <sup>3</sup>Pro - <sup>4</sup>[7-"I-Trp] - <sup>5</sup>Ile - <sup>6</sup>Leu - OH
- (13) [<sup>1</sup>M-DTPA] - <sup>1</sup>Arg - <sup>2</sup>Arg - <sup>3</sup>Pro - <sup>4</sup>Tyr - <sup>5</sup>Ile - <sup>6</sup>Leu - OH
- (14) [<sup>1</sup>M-EDTA] - <sup>1</sup>Arg - <sup>2</sup>Arg - <sup>3</sup>Pro - <sup>4</sup>Tyr - <sup>5</sup>Ile - <sup>6</sup>Leu - OH
- (15) [<sup>1</sup>M-(2-iminothiolane)] - <sup>1</sup>Arg - <sup>2</sup>Arg - <sup>3</sup>Pro - <sup>4</sup>Tyr - <sup>5</sup>Ile - <sup>6</sup>Leu - OH
- (16) [<sup>1</sup>M-(4-mercaptomethyl-2-iminothiolane)] - <sup>1</sup>Arg - <sup>2</sup>Arg - <sup>3</sup>Pro - <sup>4</sup>Tyr - <sup>5</sup>Ile - <sup>6</sup>Leu - OH ;

wherein "I is <sup>123</sup>I, <sup>125</sup>I or <sup>131</sup>I, and "M is <sup>111</sup>In, <sup>99m</sup>Tc, <sup>186</sup>Re, <sup>166</sup>Ho, <sup>153</sup>Sm or <sup>161</sup>Tb.

The above labelled peptide compounds have been tested in a number of suitable model experiments that are predictive for in vivo application. These experiments are described in the Examples. From the results of these experiments it will be evident, that the labelled peptide compounds of the present invention have properties which make them suitable for diagnostic and therapeutic purposes. If labelled with a suitable isotope for diagnostic purposes, the peptide compound remains sufficiently long intact after administration to permit imaging of the target organ or tissue without presenting a disturbing background, for example, due to detached label. If labelled with a suitable radioisotope for therapy, such-labelled peptides are promising therapeutic agents for the treatment of a number of malignant tumours that are related to neurotensin binding places, such as small cell lung carcinoma, colon carcinoma and meningiomas.


The new radiohalogenated peptide compounds of the invention can be prepared in a manner known per se for related compounds. So the invention also relates to a method of preparing a radiohalogenated peptide compound as defined hereinbefore,  
 5 characterized in that a compound of the general formula



wherein Xaa, Xbb, Xcc, Xdd, Xee,  $n$  and  $R_1$  have the meanings given above, and which compound is substituted with a non-radioactive bromine or iodine atom, attached to  $^4\text{Tyr}$  in the 2-position of the phenyl ring, to  $^4\text{Trp}$ , or to the aryl group of substituent  $R_1$ ;  
 10 is reacted with a water-soluble halogenide, selected from  $^{123}\text{I}^-$ ,  $^{124}\text{I}^-$ ,  $^{125}\text{I}^-$ ,  $^{131}\text{I}^-$ ,  $^{76}\text{Br}^-$ ,  $^{77}\text{Br}^-$  and  $^{82}\text{Br}^-$ , in the presence of copper(I) ions, a water-soluble acid and a reducing agent.

Such a halogen exchange reaction is described in European patent no. 165630. An example of a suitable water-soluble acid is citric acid; examples of suitable reducing agents are  $\text{Sn(II)}$  salts, gentisic acid, isoascorbic acid, a monosaccharide and a sulphite.  
 20

The new metal-labelled peptide compounds of the invention can also be prepared in a manner known per se for related compounds. For this purpose the peptide molecule is derivatized with the desired chelating agent as defined hereinbefore, e.g. EDTA, DTPA, etc., directly or after introduction of a spacing group as defined above, after which the compound obtained, having the general formula  
 25



wherein:

Xaa, Xbb, Xcc, Xdd, Xee and  $n$  have the meanings given above, and  $R_2$  is a chelating group attached by an amide bond or through a spacing group to the peptide molecule;  
 35 is reacted with a metal isotope, as defined hereinbefore, in the form of a salt or of a chelate bonded to a comparatively weak chelator, in order to form a complex.



Suitable examples of salts or chelates of the desired metal isotope are:  $^{111}\text{In}$ -oxinate,  $^{99\text{m}}\text{Tc}$ -tartrate, etc. The complex-forming reaction can generally be carried out in a simple manner and under conditions that are not detrimental to the peptide.

The invention further relates to a pharmaceutical composition, comprising in addition to a pharmaceutically acceptable carrier material and, if desired, at least one pharmaceutically acceptable adjuvant, as the active substance a labelled peptide compound as defined hereinbefore.

The invention also relates to a method for detecting and locating tissues having neurotensin receptors in the body of a warm-blooded living being, which comprises (i) administering to said being a composition comprising, in a quantity sufficient for external imaging, a labelled peptide compound as defined hereinbefore, wherein said peptide is labelled with (a) a metal isotope selected from the group consisting of  $^{99\text{m}}\text{Tc}$ ,  $^{203}\text{Pb}$ ,  $^{67}\text{Ga}$ ,  $^{68}\text{Ga}$ ,  $^{72}\text{As}$ ,  $^{111}\text{In}$ ,  $^{113\text{m}}\text{In}$ ,  $^{97}\text{Ru}$ ,  $^{62}\text{Cu}$ ,  $^{64}\text{Cu}$ ,  $^{52}\text{Fe}$ ,  $^{52\text{m}}\text{Mn}$ ,  $^{51}\text{Cr}$ , or (b) with a radioactive halogen isotope, selected from  $^{123}\text{I}$ ,  $^{131}\text{I}$ ,  $^{75}\text{Br}$ ,  $^{76}\text{Br}$  and  $^{77}\text{Br}$ , and thereupon (ii) subjecting said being to external imaging to determine the targeted sites in the body of said being in relation to the background activity.

The invention also relates to a method of intraoperatively detecting and locating tissues having neurotensin receptors in the body of a warm-blooded living being, which comprises (1) administering to said being a composition comprising, in a quantity sufficient for detection by a gamma detecting probe, a peptide compound as defined above, labelled with a radioisotope, selected from  $^{161}\text{Tb}$ ,  $^{123}\text{I}$  and  $^{125}\text{I}$ , preferably  $^{161}\text{Tb}$  or  $^{125}\text{I}$ , and thereupon (ii), after allowing the active substance to be taken up in said tissues and after blood clearance of radioactivity, subjecting said being to a radioimmunodetection technique in the relevant area of the body of said being, by using a gamma detecting probe.

The above radioisotope, viz. in particular  $^{161}\text{Tb}$  or  $^{125}\text{I}$ , allows the use of a such-labelled peptide compound in the technique of radioguided surgery, wherein relevant tissues in the body of a patient can be

detected and located intraoperatively by means of a gamma detecting probe. The surgeon can, intraoperatively, use this probe to find the lesions in which uptake of the compound labelled with said radioisotope, which is a low-energy gamma photon emitter, has taken place.

As mentioned hereinbefore, it is known from literature that certain tumour cells, such as small cell lung carcinoma, colon carcinoma and meningiomas, have neurotensin receptors on their surface. Therefore, the peptide compounds of the present invention, provided they are radiolabelled with isotopes suitable for the purpose, can be used for the therapeutic treatment of these tumours. So the invention further relates to a method for the therapeutic treatment of tumours, having on their surface a high density of neurotensin receptor sites, in the body of a warm-blooded living being, which comprises administering to said being a composition comprising, in a quantity effective for combating or controlling tumours, a peptide compound as defined above, said peptide compound being labelled with a metal isotope selected from the group consisting of  $^{186}\text{Re}$ ,  $^{188}\text{Re}$ ,  $^{77}\text{As}$ ,  $^{90}\text{Y}$ ,  $^{67}\text{Cu}$ ,  $^{169}\text{Er}$ ,  $^{121}\text{Sn}$ ,  $^{127}\text{Te}$ ,  $^{142}\text{Pr}$ ,  $^{140}\text{Pr}$ ,  $^{198}\text{Au}$ ,  $^{199}\text{Au}$ ,  $^{161}\text{Tb}$ ,  $^{109}\text{Pd}$ ,  $^{165}\text{Dy}$ ,  $^{149}\text{Pm}$ ,  $^{151}\text{Pm}$ ,  $^{153}\text{Sm}$ ,  $^{157}\text{Gd}$ ,  $^{159}\text{Gd}$ ,  $^{166}\text{Ho}$ ,  $^{172}\text{Tm}$ ,  $^{169}\text{Yb}$ ,  $^{175}\text{Yb}$ ,  $^{177}\text{Lu}$ ,  $^{105}\text{Rh}$  and  $^{111}\text{Ag}$ .

In case a radioactive labelled peptide compound is used as a diagnostic agent, it is frequently impossible to put the ready-for-use composition at the disposal of the user, in connection with the often poor shelf life of the radiolabelled compound and/or the short half-life of the radionuclide used. In such cases the user will carry out the labelling reaction with the radionuclide in the clinical hospital or laboratory. For this purpose the various reaction ingredients are then offered to the user in the form of a so-called "kit". It will be obvious that the manipulations necessary to perform the desired reaction should be as simple as possible to enable the user to prepare from the kit the radioactive labelled composition by using the facilities that are at his disposal. Therefore the invention also relates to a kit for preparing a radiopharmaceutical composition.

Such a kit according to the present invention may comprise (i) a peptide

compound having a selective affinity to neurotensin receptors and having the general formula II as presented hereinbefore, to which substance, if desired, an inert pharmaceutically acceptable carrier and/or formulating agents and/or adjuvants is/are added, (ii) a solution of a salt or chelate of a suitable radionuclide, and (iii) instructions for use with a prescription for reacting the ingredients present in the kit.

Suitable radionuclides for the above kit are the following metal isotopes:  $^{203}\text{Pb}$ ,  $^{67}\text{Ga}$ ,  $^{68}\text{Ga}$ ,  $^{72}\text{As}$ ,  $^{111}\text{In}$ ,  $^{113\text{m}}\text{In}$ ,  $^{97}\text{Ru}$ ,  $^{62}\text{Cu}$ ,  $^{99\text{m}}\text{Tc}$ ,  $^{186}\text{Re}$ ,  $^{188}\text{Re}$ ,  $^{64}\text{Cu}$ ,  $^{52}\text{Fe}$ ,  $^{52\text{m}}\text{Mn}$ ,  $^{51}\text{Cr}$ ,  $^{77}\text{As}$ ,  $^{90}\text{Y}$ ,  $^{67}\text{Cu}$ ,  $^{169}\text{Er}$ ,  $^{121}\text{Sn}$ ,  $^{127}\text{Te}$ ,  $^{142}\text{Pr}$ ,  $^{143}\text{Pr}$ ,  $^{198}\text{Au}$ ,  $^{199}\text{Au}$ ,  $^{161}\text{Tb}$ ,  $^{109}\text{Pd}$ ,  $^{165}\text{Dy}$ ,  $^{149}\text{Pm}$ ,  $^{151}\text{Pm}$ ,  $^{153}\text{Sm}$ ,  $^{157}\text{Gd}$ ,  $^{159}\text{Gd}$ ,  $^{166}\text{Ho}$ ,  $^{172}\text{Tm}$ ,  $^{169}\text{Yb}$ ,  $^{175}\text{Yb}$ ,  $^{177}\text{Lu}$ ,  $^{105}\text{Rh}$  and  $^{111}\text{Ag}$ .

Preferably the peptide compound to be used as an ingredient of the above kit has been modified by a reaction with a chelating agent as defined hereinbefore. The resulting peptide conjugate provides a facility for firmly attaching the radionuclide in a simple manner. Suitable chelating agents for modifying the peptide are described in detail hereinbefore. N-containing di- or polyacetic acids or their derivatives, such as the compounds mentioned before, have proved to be pre-eminently suitable for attaching various metal radionuclides, such as In-111 and In-113m, to the peptide molecules. The kit to be supplied to the user may also comprise the ingredient(s) defined sub (i) above, together with instructions for use, whereas the solution of a salt or chelate of the radionuclide, defined sub (ii) above, which solution has a limited shelf life, may be put to the disposal of the user separately.

In case the kit serves to prepare a radiopharmaceutical composition labelled with Tc-99m, Re-186 or Re-188, such a kit according to the present invention may comprise, in addition to the ingredient(s) defined sub (i) above, (ii) a reducing agent and, if desired, a chelator, and (iii) instructions for use with a prescription for reacting the ingredients of the kit with Tc-99m in the form of a pertechnetate solution, or with Re-186 or Re-188 in the form of a perrhenate solution. If desired, the ingredients of the kit may be combined, provided they are compatible. The kit should comprise a reducing agent to reduce the pertechnetate or perrhenate, for example, a dithionite, a metallic reducing agent or a complex-stabilizing reducing agent, e.g.  $\text{SnCl}_2$ ,  $\text{Sn(II)-tartrate}$ ,  $\text{Sn(II)-phosphonate}$  or -pyrophosphate, or  $\text{Sn(II)-}$

glucoheptonate. The pertechnetate or perrhenate solution can simply be obtained by the user from a suitable generator.

5 In a preferred embodiment the kit according to the present invention comprises a modified peptide or a peptide conjugate, obtained by modifying the peptide as defined hereinbefore by a treatment with a chelating agent. Suitable chelating agents have been described hereinbefore.

10 When the radionuclide is present in the kit itself, the complex forming reaction with the peptide conjugate can simply be produced by combining the components in a neutral medium and causing them to react. For that purpose the radionuclide may be presented to the peptide conjugate in the form of a chelate bonded to a comparatively weak chelator, as  
15 described hereinbefore.

When the kit comprises a peptide conjugate as defined hereinbefore and is intended for the preparation of a radiopharmaceutical composition, labelled with Tc-99m, Re-186 or Re-188, the radionuclide will preferably  
20 be added separately in the form of a pertechnetate or perrhenate solution. In that case the kit will comprise a suitable reducing agent and, if desired, a chelator, the former to reduce the pertechnetate or the perrhenate. As a reducing agent may be used, for example, a dithionite or a metallic reducing agent. The ingredients may optionally  
25 be combined, provided they are compatible. Such a monocomponent kit, in which the combined ingredients are preferably lyophilized, is excellently suitable for being reacted, by the user, with the radionuclide solution. As a reducing agent for the above-mentioned kits is preferably used a metallic reducing agent, for example, Sn(II),  
30 Ce(III), Fe(II), Cu(I), Ti(III) or Sb(III); Sn(II) is excellently suitable. The peptide constituent of the above-mentioned kits, i.e. preferably the peptide conjugate, may be supplied as a solution, for example, in the form of a physiological saline solution, or in some buffer solution, but is preferably present in a dry condition, for  
35 example, in the lyophilized condition. When used as a component for an injection liquid it should be sterile, in which, when the constituent is in the dry state, the user should preferably use a sterile physiological

saline solution as a solvent. If desired, the above-mentioned constituent may be stabilized in the conventional manner with suitable stabilizers, for example, ascorbic acid, gentisic acid or salts of these acids, or it may comprise other auxiliary agents, for example, fillers, such as glucose, lactose, mannitol, and the like.

The invention will now be described in greater detail with reference to the following specific Examples.

#### Example I

##### Synthesis of radiolabelled peptide compounds (1), (2), (3) and (4)

(a) Synthesis of the corresponding bromo-substituted peptide compounds. The starting bromo-substituted peptides, viz. bromo-substituted benzoyl-Arg-Arg-Pro-Tyr-Ile-Leu-OH and phenylacetyl-Arg-Arg-Pro-Tyr-Ile-Leu-OH, are synthesized by using the solid phase peptide synthesis (SPPS) methodology, employing N- $\alpha$ -tert.-butyloxycarbonyl (Boc) protected amino acids. During this synthesis the side chain of Arg is protected with a tosyl (Tos) group, the phenolic side chain of tyrosine with o-bromobenzyloxycarbonyl (o-Br-Z) group.

The first amino acid is attached to a resin (chloromethylpolystyrene, crosslinked with 1% of 1,4-divinylbenzene) via a Cs-salt of the desired amino acid, i.e. leucine. The substitution grade (0.60 mmol/g resin) is determined via the picric acid method.

The peptide is assembled using a repetitive cycle, as follows:

- deprotection of the main-chain protecting group, i.e. the Boc group, with a trifluoroacetic acid (TFA) solution (TFA/dichloromethane/anisole, 49/49/2);
- neutralisation of the TFA salt with diisopropylethylamine (DIEA);
- coupling of the free amine with the carboxylic function of the next amino acid; N,N'-dicyclohexylcarbodiimide (DCC) and 1-hydroxybenzotriazole (HOBt) are used as coupling agents.

This cycle is repeated until the peptide chain is assembled.

The peptide is cleaved from the resin by reaction with liquid HF. Acetic acid extraction is used to separate the peptide from the resin.

After lyophilization, the crude peptide is purified by RP-HPLC, using acetonitrile/H<sub>2</sub>O/TFA - x/(100-x)/0.001 with a gradient of x = 15%  $\rightarrow$  40% (0 - 20 min.) and a Vydac<sup>®</sup> RP-C<sub>18</sub> column.

Finally the purity of the peptide is verified by analytical RP-HPLC and thin layer chromatography. The peptide is characterized by FAB mass spectrometry. The four bromo-substituted peptide compounds are obtained in yields varying between 31 and 61%.

(b) Radioiodination with  $^{131}\text{I}$ .

The peptides are radioiodinated by nucleophilic displacement of bromine by iodine-131, according to the procedure as described in European patent no. 165630. In a typical example:

stock solution: 1 mg  $\text{SnSO}_4$ ,

25 mg 2,5-dihydroxybenzoic acid (gentisic acid),

35 mg citric acid monohydrate,

in 2.5 ml 10%  $\text{AcOH.aq}$  (v/v);

$\text{Cu}^{2+}$ -solution: 32.5 mg  $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$  in 10 ml water (0.013 M).

The peptide (1 mg) is dissolved in 500  $\mu\text{l}$  of stock solution and 60  $\mu\text{l}$  of  $\text{Cu}^{2+}$ -solution. The solution is sonicated until all solutes are dissolved. The clear solution is flushed with  $\text{N}_2$  for 5 min.

After the addition of 10  $\mu\text{l}$   $\text{Na}^{131}\text{I}$  (132  $\mu\text{Ci}$ ), the reaction mixture is heated at  $140^\circ\text{C}$  for 1 hour. The mixture is finally cooled to ambient temperature. RP-HPLC is carried out to calculate the iodination yield: acetonitrile/ $\text{H}_2\text{O}$ /TFA -  $x/(100-x)/0.001$  with a gradient of  $x = 15\% \rightarrow 40\%$  (0 - 20 min.); Lichrospher<sup>®</sup> 100 RP-18 (5 $\mu\text{m}$ ) column; flow-rate 1 ml/min. The desired radioiodinated peptides (1), (2), (3) and (4) are obtained in yields of 84%, 54%, 86% and 55%, respectively (not optimized).

#### Example II

##### Biological experiments

Binding assay experiment.

Binding assays are performed by studying the inhibition of [ $^3\text{H}$ ]neurotensin binding to guinea pig forebrain membranes. The following results are obtained, using the unlabelled acetyl compound Ac-Arg-Arg-Pro-Tyr-Ile-Leu-OH (a) as the reference.

The affinity is expressed as the  $\text{pIC}_{50}$  value (i.e. the concentration of peptide analogue yielding 50% inhibition).

peptide	pIC <sub>50</sub> (M)
(a)	8.11
(1)	8.05
(2)	7.87
(3)	8.07
(4)	8.10

From the above figures it will be clear that the radioiodinated analogues (1) through (4) show affinities to neurotensin receptor sites in this binding assay which are comparable with that of the unlabelled compound.

#### Enzymatic stability.

The enzymatic stability of the four radioiodinated analogues (1), (2), (3) and (4) is determined in vitro in human serum in a conventional manner, showing that this stability is sufficient.

Structure-activity studies of neurotensin and its analogues reveals that the same relationship, viz. structure to activity, exists in rat brain receptors as in tumour receptors. This indicates that the receptors in rat brain are the same as those in tumour cells. Therefore the above experiments show, that the tested compounds are promising tools in diagnosing malignancies related to neurotensin receptors.

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

## (i) APPLICANT:

- (A) NAME: Mallinckrodt Medical B.V.
- (B) STREET: Westerduinweg 3
- (C) CITY: Petten
- (E) COUNTRY: The Netherlands
- (F) POSTAL CODE (ZIP): none

(ii) TITLE OF INVENTION: Labelled peptide compounds

(iii) NUMBER OF SEQUENCES: 16

## (iv) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version #1.25

(EPO)

## (v) CURRENT APPLICATION DATA:

APPLICATION NUMBER: EP 94200409.4

**SUBSTITUTE SHEET (RULE 26)**



## (2) INFORMATION FOR SEQ ID NO: 1:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 6 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

## (ix) FEATURE:

(A) NAME/KEY: misc-feature

(B) LOCATION: 1

(D) OTHER INFORMATION: Xaa is (2\*I-phenyl)acetyl-Arg

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

Xaa Arg Pro Tyr Ile Leu

1 5

SUBSTITUTE SHEET (RULE 26)

## (2) INFORMATION FOR SEQ ID NO: 2:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 6 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

## (ix) FEATURE:

(A) NAME/KEY: misc-feature

(B) LOCATION: 1

(D) OTHER INFORMATION: Xaa is (4-\*I-phenyl)acetyl-Arg

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Xaa Arg Pro Tyr Ile Leu  
1 5

**SUBSTITUTE SHEET (RULE 26)**

## (2) INFORMATION FOR SEQ ID NO: 3:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 6 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

## (ix) FEATURE:

(A) NAME/KEY: misc-feature

(B) LOCATION: 1

(D) OTHER INFORMATION: Xaa is 2-\*I-benzoyl-Arg

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

Xaa Arg Pro Tyr Ile Leu

1 5

**SUBSTITUTE SHEET (RULE 26)**

## (2) INFORMATION FOR SEQ ID NO: 4:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 6 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

## (ix) FEATURE:

(A) NAME/KEY: misc-feature

(B) LOCATION: 1

(D) OTHER INFORMATION: Xaa is 4-\*I-benzoyl-Arg

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

Xaa Arg Pro Tyr Ile Leu

1 5

**SUBSTITUTE SHEET (RULE 26)**

## (2) INFORMATION FOR SEQ ID NO: 5:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 6 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

## (ix) FEATURE:

(A) NAME/KEY: mixc-feature

(B) LOCATION: 1

(D) OTHER INFORMATION: Xaa is (4-F-2-\*I-phenyl)acetyl-Arg

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

Xaa Arg Pro Tyr Ile Leu  
1 5

**SUBSTITUTE SHEET (RULE 26)**

## (2) INFORMATION FOR SEQ ID NO: 6:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 6 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

## (ix) FEATURE:

(A) NAME/KEY: misc-feature

(B) LOCATION: 1

(D) OTHER INFORMATION: Xaa is (4-F-2-\*I-benzoyl)-Arg

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

Xaa Arg Pro Tyr Ile Leu  
1                      5

**SUBSTITUTE SHEET (RULE 26)**

## (2) INFORMATION FOR SEQ ID NO: 7:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 6 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

## (ix) FEATURE:

(A) NAME/KEY: misc-feature

(B) LOCATION: 1

(D) OTHER INFORMATION: Xaa is (2-\*I-phenyl)acetyl-Lys

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

Xaa Arg Pro Tyr Ile Leu  
1                   5

**SUBSTITUTE SHEET (RULE 26)**

## (2) INFORMATION FOR SEQ ID NO: 8:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 6 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

## (ix) FEATURE:

(A) NAME/KEY: misc-feature

(B) LOCATION: 1

(D) OTHER INFORMATION: Xaa is (2-\*I-phenyl)acetyl-Arg

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

Xaa Lys Pro Tyr Ile Leu

1 5

SUBSTITUTE SHEET (RULE 26)



## (2) INFORMATION FOR SEQ ID NO: 9:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 6 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

## (ix) FEATURE:

(A) NAME/KEY: misc-feature

(B) LOCATION: 1

(D) OTHER INFORMATION: Xaa is Ac-Arg

## (ix) FEATURE:

(A) NAME/KEY: misc-feature

(B) LOCATION: 4

(D) OTHER INFORMATION: Xaa is 2-\*I-Tyr

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

Xaa Arg Pro Xaa Ile Leu  
1 5

**SUBSTITUTE SHEET (RULE 26)**

## (2) INFORMATION FOR SEQ ID NO: 10:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 6 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

## (ix) FEATURE:

(A) NAME/KEY: misc-feature

(B) LOCATION: 1

(D) OTHER INFORMATION: Xaa is Ac-Arg

## (ix) FEATURE:

(A) NAME/KEY: misc-feature

(B) LOCATION: 4

(D) OTHER INFORMATION: Xaa is 2-\*I-Trp

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

Xaa Arg Pro Xaa Ile Leu

1

5

SUBSTITUTE SHEET (RULE 26)

## (2) INFORMATION FOR SEQ ID NO: 11:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 6 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

## (ix) FEATURE:

(A) NAME/KEY: misc-feature

(B) LOCATION: 1

(D) OTHER INFORMATION: Xaa is Ac-Arg

## (ix) FEATURE:

(A) NAME/KEY: misc-feature

(B) LOCATION: 4

(D) OTHER INFORMATION: Xaa is 5-\*I-Trp

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

Xaa Arg Pro Xaa Ile Leu

1

5

SUBSTITUTE SHEET (RULE 26)

## (2) INFORMATION FOR SEQ ID NO: 12:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 6 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: peptide
- (iii) HYPOTHETICAL: NO
- (iii) ANTI-SENSE: NO
- (ix) FEATURE:
  - (A) NAME/KEY: misc-feature
  - (B) LOCATION: 1
  - (D) OTHER INFORMATION: Xaa is Ac-Arg
- (ix) FEATURE:
  - (A) NAME/KEY: misc-feature
  - (B) LOCATION: 4
  - (D) OTHER INFORMATION: Xaa is 7-<sup>\*</sup>I-Trp
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:  

Xaa	Arg	Pro	Xaa	Ile	Leu
1				5	

SUBSTITUTE SHEET (RULE 26)

## (2) INFORMATION FOR SEQ ID NO: 13:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 6 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

## (ix) FEATURE:

(A) NAME/KEY: misc-feature

(B) LOCATION: 1

(D) OTHER INFORMATION: Xaa =[\*M-DTPA]-Arg

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

Xaa Arg Pro Tyr Ile Leu  
1           5

**SUBSTITUTE SHEET (RULE 26)**

## (2) INFORMATION FOR SEQ ID NO: 14:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 6 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

## (ix) FEATURE:

- (A) NAME/KEY: misc-feature
- (B) LOCATION: 1
- (D) OTHER INFORMATION: Xaa is [\*M-EDTA]-Arg

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

Xaa Arg Pro Tyr Ile Leu  
1           5

**SUBSTITUTE SHEET (RULE 26)**

## (2) INFORMATION FOR SEQ ID NO: 15:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 6 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

## (ix) FEATURE:

(A) NAME/KEY: misc-feature

(B) LOCATION: 1

(D) OTHER INFORMATION: Xaa is [\*M-(2-iminothiolane)]-Arg

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

Xaa Arg Pro Tyr Ile Leu  
1 5

**SUBSTITUTE SHEET (RULE 26)**

## (2) INFORMATION FOR SEQ ID NO: 16:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 6 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

## (ix) FEATURE:

(A) NAME/KEY: misc-feature

(B) LOCATION: 1

(D) OTHER INFORMATION: Xaa is [<sup>\*</sup>M-(4-mercaptomethyl-2-iminothiolane)]-Arg

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:


Xaa Arg Pro Tyr Ile Leu  
1                      5

SUBSTITUTE SHEET (RULE 26)



Claims

1. A labelled peptide compound, wherein the peptide has a selective neurotensin receptor affinity and is represented by the general formula



wherein:

$R_1$  is a ( $C_1$ - $C_3$ )alkanoyl group, an arylcarbonyl group, an aryl- ( $C_1$ - $C_3$ )alkanoyl group, or a chelating group attached by an amide bond or through a spacing group to the peptide molecule;

Xaa and Xbb are each individually Arg or Lys;

Xcc is an unsubstituted or substituted cyclic amino acid, preferably selected from Pro and Hyp;

Xdd is Tyr, Trp or Phe;

Xee is Leu, Ile or t.-butylalanine; and

$n$  is 0 or 1;

and wherein said peptide is labelled with (a) a metal isotope selected from the group consisting of  $^{99m}\text{Tc}$ ,  $^{203}\text{Pb}$ ,  $^{67}\text{Ga}$ ,  $^{68}\text{Ga}$ ,  $^{72}\text{As}$ ,  $^{111}\text{In}$ ,  $^{113m}\text{In}$ ,  $^{97}\text{Ru}$ ,  $^{62}\text{Cu}$ ,  $^{64}\text{Cu}$ ,  $^{52}\text{Fe}$ ,  $^{52m}\text{Mn}$ ,  $^{51}\text{Cr}$ ,  $^{186}\text{Re}$ ,  $^{188}\text{Re}$ ,  $^{77}\text{As}$ ,  $^{90}\text{Y}$ ,  $^{67}\text{Cu}$ ,  $^{169}\text{Er}$ ,  $^{121}\text{Sn}$ ,  $^{127}\text{Te}$ ,  $^{142}\text{Pr}$ ,  $^{143}\text{Pr}$ ,  $^{198}\text{Au}$ ,  $^{199}\text{Au}$ ,  $^{161}\text{Tb}$ ,  $^{109}\text{Pd}$ ,  $^{163}\text{Dy}$ ,  $^{149}\text{Pm}$ ,  $^{151}\text{Pm}$ ,  $^{153}\text{Sm}$ ,  $^{157}\text{Gd}$ ,  $^{159}\text{Gd}$ ,  $^{166}\text{Ho}$ ,  $^{172}\text{Tm}$ ,  $^{169}\text{Yb}$ ,  $^{175}\text{Yb}$ ,  $^{177}\text{Lu}$ ,  $^{105}\text{Rh}$  and  $^{111}\text{Ag}$ , or

(b) with a radioactive halogen isotope;

on the understanding that:


(i) if the label is a metal isotope,  $R_1$  represents a chelating group for chelating said isotope; and

(ii) if the label is a radioactive halogen isotope, said halogen isotope is attached to  $^4\text{Tyr}$  in the 2-position of the phenyl ring, to  $^4\text{Trp}$ , or to the aryl group of substituent  $R_1$ .

2. A labelled peptide compound as claimed in claim 1, wherein said peptide is labelled with a radioactive halogen isotope selected from the group consisting of  $^{123}\text{I}$ ,  $^{124}\text{I}$ ,  $^{125}\text{I}$ ,  $^{131}\text{I}$ ,  $^{75}\text{Br}$ ,  $^{76}\text{Br}$ ,  $^{77}\text{Br}$  and  $^{82}\text{Br}$ .

3. A labelled peptide compound as claimed in claim 1, wherein said peptide is labelled with a metal isotope, chelated by a chelating group

$R_1$  derived from ethylene diamine tetra-acetic acid (EDTA), diethylene triamine penta-acetic acid (DTPA), cyclohexyl 1,2-diamine tetra-acetic acid (CDTA), ethyleneglycol-0,0'-bis(2-aminoethyl)-N,N,N',N'-tetra-acetic acid (EGTA), N,N-bis(hydroxybenzyl)-ethylenediamine-N,N'-diacetic acid (HBED), triethylene tetramine hexa-acetic acid (TTHA), 1,4,7,10-tetraazacyclododecane-N,N',N'',N'''-tetra-acetic acid (DOTA), hydroxyethyl diamine triacetic acid (HEDTA), 1,4,8,11-tetra-azacyclo-tetradecane-N,N',N'',N'''-tetra-acetic acid (TETA), substituted DTPA, substituted EDTA, or from a compound of the general formula




(III)

wherein R is a branched or non-branched, optionally substituted hydrocarbyl radical, which may be interrupted by one or more hetero-atoms selected from N, O and S and/or by one or more NH groups, and

Y is a group which is capable of reacting with an amino group of the peptide and which is preferably selected from the group consisting of carbonyl, carbimidoyl, N-(C<sub>1</sub>-C<sub>6</sub>)alkylcar-bimidoyl, N-hydroxycarbimidoyl and N-(C<sub>1</sub>-C<sub>6</sub>)alkoxycar-bimidoyl;

and


wherein said optionally present spacing group has the general formula



wherein R<sub>3</sub> is a C<sub>1</sub>-C<sub>10</sub> alkylene group, a C<sub>1</sub>-C<sub>10</sub> alkylidene group or a C<sub>2</sub>-C<sub>10</sub> alkenylene group, and X is a thiocarbonyl group or a methylcarbonyl group.

4. A method of preparing a radiohalogenated peptide compound as claimed in claim 2,

characterized in that a compound of the general formula




wherein Xaa, Xbb, Xcc, Xdd, Xee,  $n$  and  $R_1$  have the meanings given in claim 1, and

which compound is substituted with a non-radioactive bromine or iodine atom, attached to  $^4\text{Tyr}$  in the 2-position of the phenyl ring, to  $^4\text{Trp}$ , or to the aryl group of substituent  $R_1$ ;

is reacted with a water-soluble halogenide, selected from  $^{123}\text{I}^-$ ,  $^{124}\text{I}^-$ ,  $^{125}\text{I}^-$ ,  $^{131}\text{I}^-$ ,  $^{75}\text{Br}^-$ ,  $^{76}\text{Br}^-$ ,  $^{77}\text{Br}^-$  and  $^{82}\text{Br}^-$ , in the presence of copper(I) ions, a water-soluble acid and a reducing agent.

5. A method of preparing a metal-labelled peptide compound as claimed in claim 3,

characterized in that a compound of the general formula



wherein:

Xaa, Xbb, Xcc, Xdd, Xee and  $n$  have the meaning given in claim 1, and

$R_2$  is a chelating group attached by an amide bond or through a spacing group to the peptide molecule;

is reacted with a metal isotope as defined in claim 1 in the form of a salt or of a chelate bonded to a comparatively weak chelator, in order to form a complex.

6. A pharmaceutical composition, comprising in addition to a pharmaceutically acceptable carrier material and, if desired, at least one pharmaceutically acceptable adjuvant, as the active substance a labelled peptide compound as claimed in claim 1, 2 or 3.

7. A method for detecting and locating tissues having neurotensin receptors in the body of a warm-blooded living being, which comprises (i) administering to said being a composition comprising, in a quantity sufficient for external imaging, a labelled peptide compound as claimed in claim 1, 2 or 3, wherein said peptide is labelled with (a) a metal isotope selected from the group consisting of  $^{99\text{m}}\text{Tc}$ ,  $^{203}\text{Pb}$ ,  $^{67}\text{Ga}$ ,  $^{68}\text{Ga}$ ,  $^{72}\text{As}$ ,  $^{111}\text{In}$ ,  $^{113\text{m}}\text{In}$ ,  $^{97}\text{Ru}$ ,  $^{62}\text{Cu}$ ,  $^{64}\text{Cu}$ ,  $^{52}\text{Fe}$ ,  $^{52\text{m}}\text{Mn}$ ,  $^{51}\text{Cr}$ , or (b) with a radioactive

halogen isotope, selected from  $^{123}\text{I}$ ,  $^{131}\text{I}$ ,  $^{75}\text{Br}$ ,  $^{76}\text{Br}$  and  $^{77}\text{Br}$ , and thereupon  
 (ii) subjecting said being to external imaging to determine the targeted  
 sites in the body of said being in relation to the background activity.

5 8. A method of intraoperatively detecting and locating tissues having  
 neurotensin receptors in the body of a warm-blooded living being, which  
 comprises (1) administering to said being a composition comprising, in  
 a quantity sufficient for detection by a gamma detecting probe, a  
 10 labelled peptide compound as claimed in claim 1, wherein the  
 radioisotope is  $^{161}\text{Tb}$ ,  $^{123}\text{I}$  or  $^{125}\text{I}$ , preferably  $^{161}\text{Tb}$  or  $^{125}\text{I}$ , and thereupon  
 (ii), after allowing the active substance to be taken up in said tissues  
 and after blood clearance of radioactivity, subjecting said being to a  
 radioimmunodetection technique in the relevant area of the body of said  
 15 being, by using a gamma detecting probe.

9. A method for the therapeutic treatment of tumours, having on their  
 surface a high density of neurotensin receptor sites, in the body of a  
 warm-blooded living being, which comprises administering to said being  
 a composition comprising, in a quantity effective for combating or  
 20 controlling tumours, a peptide compound as claimed in claim 3, said  
 peptide compound being labelled with a metal isotope selected from the  
 group consisting of  $^{186}\text{Re}$ ,  $^{188}\text{Re}$ ,  $^{77}\text{As}$ ,  $^{90}\text{Y}$ ,  $^{67}\text{Cu}$ ,  $^{169}\text{Er}$ ,  $^{121}\text{Sn}$ ,  $^{127}\text{Te}$ ,  $^{142}\text{Pr}$ ,  $^{143}\text{Pr}$ ,  
 $^{198}\text{Au}$ ,  $^{199}\text{Au}$ ,  $^{161}\text{Tb}$ ,  $^{109}\text{Pd}$ ,  $^{165}\text{Dy}$ ,  $^{149}\text{Pm}$ ,  $^{151}\text{Pm}$ ,  $^{153}\text{Sm}$ ,  $^{157}\text{Gd}$ ,  $^{159}\text{Gd}$ ,  $^{166}\text{Ho}$ ,  $^{172}\text{Tm}$ ,  $^{169}\text{Yb}$ ,  
 25  $^{175}\text{Yb}$ ,  $^{177}\text{Lu}$ ,  $^{105}\text{Rh}$  and  $^{111}\text{Ag}$ .

10. A kit for preparing a radiopharmaceutical composition, comprising  
 (i) a peptide compound having a selective affinity to neurotensin  
 receptors and having the general formula II, to which compound, if  
 30 desired, an inert pharmaceutically acceptable carrier and/or formulating  
 agents and/or adjuvants is/are added, (ii) a solution of a salt or  
 chelate of a metal isotope selected from the group consisting of  $^{203}\text{Pb}$ ,  
 $^{67}\text{Ga}$ ,  $^{68}\text{Ga}$ ,  $^{72}\text{As}$ ,  $^{111}\text{In}$ ,  $^{113\text{m}}\text{In}$ ,  $^{97}\text{Ru}$ ,  $^{62}\text{Cu}$ ,  $^{99\text{m}}\text{Tc}$ ,  $^{186}\text{Re}$ ,  $^{188}\text{Re}$ ,  $^{64}\text{Cu}$ ,  $^{52}\text{Fe}$ ,  $^{52\text{m}}\text{Mn}$ ,  
 $^{51}\text{Cr}$ ,  $^{77}\text{As}$ ,  $^{90}\text{Y}$ ,  $^{67}\text{Cu}$ ,  $^{169}\text{Er}$ ,  $^{121}\text{Sn}$ ,  $^{127}\text{Te}$ ,  $^{142}\text{Pr}$ ,  $^{143}\text{Pr}$ ,  $^{198}\text{Au}$ ,  $^{199}\text{Au}$ ,  $^{161}\text{Tb}$ ,  $^{109}\text{Pd}$ ,  
 35  $^{165}\text{Dy}$ ,  $^{149}\text{Pm}$ ,  $^{151}\text{Pm}$ ,  $^{153}\text{Sm}$ ,  $^{157}\text{Gd}$ ,  $^{159}\text{Gd}$ ,  $^{166}\text{Ho}$ ,  $^{172}\text{Tm}$ ,  $^{169}\text{Yb}$ ,  $^{175}\text{Yb}$ ,  $^{177}\text{Lu}$ ,  $^{105}\text{Rh}$  and  
 $^{111}\text{Ag}$ , and (iii) instructions for use with a prescription for reacting the  
 ingredients present in the kit.

11. A kit for preparing a radiopharmaceutical composition, comprising  
(i) a peptide compound having a selective affinity to neurotensin  
receptors and having the general formula II, to which compound, if  
desired, an inert pharmaceutically acceptable carrier and/or formulating  
agents and/or adjuvants is/are added, (ii) a reducing agent, and, if  
desired, a chelator, said ingredients (i) and (ii) optionally being  
combined, and (iii) instructions for use with a prescription for  
reacting the ingredients of the kit with  $^{99m}\text{Tc}$  in the form of a  
pertechnetate solution or with  $^{186}\text{Re}$  or  $^{188}\text{Re}$  in the form of a perrhenate  
solution.

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US95/02131

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A61K 38/00, 49/00; C07K 5/00, 7/00, 17/00

US CL : 424/1.41, 1.45, 1.69; 514/16; 530/328

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/1.41, 1.45, 1.69; 514/16; 530/328

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

USPTO APS

search terms: neurotensin, radiolabel

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Biochemical and Biophysical Research Communications, Volume 120, No. 3, issued 16 May 1984, J. Sadoul et al, "Preparation of Neurotensin Selectively Iodinated on the Tyrosine 3 Residue. Biological Activity and Binding Properties on Mammalian Neurotensin Receptors", pages 812-819.	1-11

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

*A* document defining the general state of the art which is not considered to be of particular relevance	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*E* earlier document published on or after the international filing date	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*O* document referring to an oral disclosure, use, exhibition or other means	*G* document member of the same patent family
*P* document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

19 APRIL 1995

Date of mailing of the international search report

15 MAY 1995

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

SHEELA J. HUFF

Telephone No. (703) 308-0196

Form PCT/ISA/210 (second sheet)(July 1992)\*